

RX J0911+05: A Massive Cluster Lens at $z=0.769$

Jean-Paul Kneib¹

Observatoire Midi-Pyrénées, 14 Av. E.Belin, 31400 Toulouse, France
 kneib@ast.obs-mip.fr

Judith G. Cohen

Palomar Observatory, Mail Stop 105-24, California Institute of Technology, Pasadena, CA 91125, USA
 jlc@astro.caltech.edu

Jens Hjorth

Astronomical Observatory, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark
 jens@astro.ku.dk

ABSTRACT

We report the detection of a massive high-redshift cluster of galaxies near the quadruple quasar RX J0911+05, using the LRIS instrument on the Keck-II telescope. The cluster is found to have a mean redshift of $\bar{z} = 0.7689 \pm 0.002$ and a velocity dispersion of $\sigma = 836^{+180}_{-200}$ km s⁻¹, based on redshift measurements for 24 member galaxies. This massive high-redshift cluster is the origin of the unusually large external shear required by lensing models of the quadruple quasar system. We predict the expected time delay depending on the exact contribution of the cluster. A measurement of the time delay and further deep lensing and X-ray observations will unravel useful properties of this serendipitously discovered high-redshift cluster, and may put interesting cosmological constraints on H_0 .

Subject headings: galaxies: clusters: general — gravitational lensing — galaxies: clusters: individual (RX J0911.4+0551)

1. Introduction

Multiply imaged quasars are among the most promising objects for measuring distances and hence cosmological parameters (Refsdal 1964; Blandford & Narayan 1992). That promise is finally starting to be realized after time delays have been measured in several systems in recent years (e.g., Schechter et al. 1997; Kundić et al. 1997c; Hjorth et al. 1999). Contrary to local estimates of the Hubble parameter, multiply imaged quasars

with time delays can be used to measure the Hubble parameter on a *cosmological* scale and hence are insensitive to local bulk flows.

Until recently, simple mass distributions have been used to model multiply imaged QSOs. However, the rapid development of observing techniques, and the availability of powerful observatories like the *Hubble Space Telescope* and the 8–10 m class telescopes have shown that most of the image configurations are more complex than allowed by simple models. As an example, the famous double QSO 0957+561, first thought to be produced by a single galaxy, is now known to be a complex system involving a massive lens plus one or more galaxy clusters (Bernstein & Fisher 1999 and references therein). In parallel to the surge in obser-

¹Visiting astronomer, of W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

vational efforts, the modeling of galaxies and their surroundings has greatly improved and is starting to provide very important insights into the distribution of dark matter in galaxies (e.g. Hjorth & Kneib 2000; Maller et al. 2000; Kochanek et al. 2000).

A major uncertainty in the mass models comes from the surroundings of the lens, in the form of “external shear”, e.g., due to either nearby galaxies, a galaxy group or a cluster close to the line of sight to the QSO (Keeton, Kochanek & Seljak 1997, Kundic et al. 1997b). It is important to underline that a reliable mass model requires not only γ , the gravitational external shear, but also the external surface mass density κ , producing the measured external shear. Therefore, a measurement of the mass distribution in the vicinity of the lensed quasar is *mandatory* in putting stringent constraints on cosmological parameters.

The quadruple lens system RX J0911+05, is potentially one of the most valuable systems for cosmological studies. It was identified by Bade et al. (1997) as a multiply imaged radio quiet QSO at $z = 2.80$, selected as an AGN candidate from the ROSAT All-Sky Survey. In high-resolution NOT and NTT images (Burud et al. 1998), RX J0911+05 was soon found to consist of four QSO components and an elongated lensing galaxy, confirmed by subsequent *HST* imaging. Regular monitoring of RX J0911+05 with the Nordic Optical Telescope is expected to give a time-delay to about 10 percent (Hjorth et al. 2000). Moreover, recently, the redshift of the main lens was measured at $z = 0.769$ with the Keck telescope (Falco, Davis & Stern, 1999, private communication). The unusual image configuration of RX J0911+05 requires a large external shear ($\gamma_{\min} = 0.15$) to be included in the lensing potential in order to reproduce the complex geometry observed in RX J0911+05. The probable source of this shear is a galaxy cluster located about $38''$ from RX J0911+05 as suggested by Burud et al. (1998). Such a cluster may contribute to the X-ray emission observed by ROSAT. The color of the presumed cluster galaxies indicates a redshift in the range of 0.6–0.8, consistent with the determined spectroscopic redshift of the main lens galaxy.

In this Letter we present the results of spectroscopic observations of RX J0911+05 conducted

at the Keck Observatory using the LRIS instrument. Section 2 summarizes the photometric and spectroscopic observations. Section 3 presents the constraints on the mass model of the high-redshift cluster of galaxies detected near the quadruple lens. A discussion of the results is presented in section 4. Throughout this paper we use a Hubble parameter $H_0 = 50h_{50}^{-1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$, which gives a physical scale of $10.37h_{50}^{-1} \text{ kpc arcsec}^{-1}$ at the cluster redshift ($z = 0.769$).

2. Observations

We observed RX J0911+05 during the December 7–8, 1999 and March 6–7, 2000 nights with the LRIS instrument (Oke et al. 1995) mounted on Keck II. Imaging was done in B, R and I and calibrated using photometric standards PG2213-006 and other stars in the SA92 field (Landolt 1992). Spectroscopic observation were conducted with the 400/8500 grating blazed at $7200 \text{ \AA}$ which gives a dispersion of $1.84 \text{ \AA/pixel}$. Standard *Figaro* and *IRAF* pipeline reduction packages were used to reduce the photometric and spectroscopic data. The Journal of these observations is given in Table 4. A color image made of the B, R and I images is presented in Figure 1.

We exposed 2 MOS masks, the first mask had 24 slitlets, and the second 30 slitlets. Object selection was limited in magnitude to $I_{AB} < 22.5$ and generally followed the color selection of Burud et al. (1998). In the first mask, a total of 25 objects were observed (there were two objects in one slitlet); 3 stars and 2 bright galaxies were used to allow a correct alignment of the mask on the objects. Similarly for the second mask, we targeted 30 galaxies, and used 3 stars for the mask alignment.

Redshifts were estimated independently by JPK and JGC, by simple line fitting techniques, as well as using the cross-correlation package *RVSAO* V2.0 in *IRAF* (Kurtz & Mink 1998). Redshift identifications are displayed on Figure 1. Table 4 summarizes the photometry and the redshift of the cluster members. Figure 2 shows the redshift histogram of the entire sample and the cluster sample. The majority of the cluster member spectra are typical of an old stellar population. Only 3 out of 23 spectra show a strong

[O II] emission line (one of them being an AGN), and 4 others show a weak [O II] line. Although our target selection was clearly biased to have a large number of early type galaxies, it clearly shows that these galaxies belong to an evolved cluster. Note that the small group of apparently redder objects detected some $10''$ SW from the lens (Burud et al. 1998) is found at the same redshift as the cluster.

3. Cluster Dynamics and Mass Model

Including the redshift of the lensing galaxy (Falco, Davis & Stern 1999, private communication) we have redshift measurements of 24 cluster members with a mean redshift $\bar{z} = 0.7689 \pm 0.002$. The velocity dispersion of the 24 galaxies is $\sigma_{\text{los}} = 836_{-200}^{+180}$ km s $^{-1}$. If one keeps only the 14 high quality spectra, we estimate the mean redshift of $\bar{z} = 0.7692 \pm 0.004$ and a velocity dispersion of $\sigma_{\text{los}} = 832_{-250}^{+175}$ km s $^{-1}$. The two estimates are in good agreement, so we will retain the first one in the following discussion. We used the biweight estimator to measure the cluster redshift and its dispersion. The errors were estimated by using the bootstrap algorithm for the mean redshift and the jackknife algorithm for the velocity dispersion (Beers, Flynn & Gebhardt 1990).

We computed the harmonic radius R_h (e.g., Nolthenius & White 1987) from our spectroscopic cluster members as

$$R_h = D_A(\bar{z}) \frac{\pi}{2} \frac{N_m(N_m - 1)}{2} (\Sigma_i \Sigma_{j>i} \theta_{ij}^{-1})^{-1}, \quad (1)$$

where θ_{ij} is the angular distance between galaxies i and j , N_m is the number of cluster members, and $D_A(\bar{z})$ is the angular diameter distance at the mean cluster redshift $\bar{z}$. The cluster virial mass can then be estimated as

$$M_V = \frac{6\sigma^2 R_h}{G}. \quad (2)$$

We found an harmonic radius of $R_h = 632h_{50}^{-1}$ kpc and a mass $M_V = 6.2_{-2.6}^{+2.9} 10^{14} M_\odot$. Assuming that the cluster follows the σ - T_X relation (e.g. Girardi et al. 1996), we derive an X-ray temperature of $T_X = 4.5 \pm 1.2$ keV.

We have thus identified a new massive cluster at high redshift. This cluster is not as massive as the cluster MS1054-03 (Tran et al. 1998; Hoekstra et al. 2000) but is similar to MS1137.5+6625 ($\sigma = 884$ km s $^{-1}$ as measured by Donahue et al. 2000).

Assuming a singular isothermal sphere centered on the brightest galaxy of the cluster, the computed velocity dispersion translates into a gravitational shear of $\gamma = \kappa = 0.11_{-0.03}^{+0.04}$ for $(\Omega, \lambda) = (1, 0)$ and $\gamma = \kappa = 0.13_{-0.03}^{+0.04}$ for $(\Omega, \lambda) = (0.3, 0.7)$ at the location of the multiple quasar. This quick estimate of the external shear is close to the value found in the modeling of the lens plus an external shear (Burud et al. 1998).

Having identified the cluster redshift and derived an estimate of the galaxy velocity dispersion, we can go a step further and try to model the system modeling mass distribution of the cluster and its galaxies in a similar way as for Abell 2218 (Kneib et al. 1996). The cluster galaxies are modeled as truncated isothermal spheres, with a velocity dispersion and truncation radius scaled with the galaxy luminosity. The center position of the cluster is set to the position of the brightest galaxy and the velocity dispersion as given by our spectroscopic survey. The constraints are the quasar positions and flux ratios (although we only give a relatively small weight to the flux ratio constraints due to possible microlensing (Burud et al. 1998), the position, orientation, and ellipticity of the main lens system as measured on the HST/NICMOS-2 image (PI: Falco). The fiducial best-fitting model found is displayed in Figure 3, where we show the surface mass density of the model. Note that the lens model favors an elongated mass distribution of the cluster aligned with the cluster light distribution. The time delay expected between images A and B is strongly dependent on the exact morphology and mass distribution of the cluster; we found that $190 < \Delta t \text{ (days)} < 260h_{50}^{-1}$. The lower time-delay value corresponds to a more massive cluster. Clearly it would be very important to further constrain the mass distribution of the cluster before trying to get a constraint on the Hubble parameter.

4. Discussion

We have identified a massive high-redshift cluster of galaxies at $\bar{z} = 0.769 \pm 0.002$ responsible for the very large external shear ($\gamma \sim 0.15$), affecting the lens potential of the quadrupole quasar RX J0911+05. The measured velocity dispersion based on 24 members leads to a velocity dispersion of $\sigma = 836_{-200}^{+180}$ km s $^{-1}$. Using these re-

sults we present a new mass model for this lens which includes the contribution of a cluster and the cluster galaxies in a similar way as Kneib et al. (1996). The predicted time delay is estimated to be $190 < \Delta t \text{ (days)} < 260h_{50}^{-1}$, depending on the exact mass distribution of the cluster.

Further observations using the WFPC2 on board HST will allow us to directly measure the predicted cluster weak shear from the distortions of faint background galaxies; X-ray observations conducted with the new X-ray satellites (*Chandra* or *XMM-Newton*) will permit to study the gas properties of this distant cluster in detail. Such complementary data will be essential in producing a reliable mass model of the system which, in combination with the foreseen measurement of the time delay, may provide the best constrained mass model for a multiple QSO, hence leading to an accurate cosmological estimate of the Hubble parameter.

This serendipitous cluster discovery is very interesting because it demonstrates that multiple quasars with large separation are efficient in revealing high-redshift collapsed structure. Indeed, there is a growing number of such multiple quasars where a cluster or group as been detected (e.g., Q0957: Bernstein & Fischer 1999; the Cloverleaf: Kneib et al. 1998a, 1998b, MG2016+116 Hattori et al. 1997; Benitez et al. 1999; Soucail et al. 2000, PG1115+080: Schechter et al. 1997, Kundic et al. 1997a). Therefore, multiple QSO system, constitute a useful way to discover and study high-redshift groups and clusters along their line of sight.

Part of the work presented is based on observations made with the NASA/ESA Hubble Space Telescope, obtained from the data archive at the Space Telescope Science Institute. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555. The entire Keck/LRIS user community owes a huge debt to Jerry Nelson, Gerry Smith, Bev Oke, and many other people who have worked to make the Keck Telescope and LRIS a reality. We are grateful to the W. M. Keck Foundation, and particularly its late president, Howard Keck, for the vision to fund the construction of the W. M. Keck Observatory.

It is a pleasure to acknowledge the efficient and

friendly support of the Keck Observatory staff. We also acknowledge useful discussions with C. Kochanek, F. Courbin, and I. Burud. This research was supported by CNRS/INSU for JPK and by the Danish Natural Science Research Council (SNF) for JH.

REFERENCES

- Bade, N., Siebert, J., Lopez, S., Voges, W. & Reimers, D. 1997, *A&A*, 317, L13
- Burud, I., Courbin, F., Lidman, C., Jaunsen, A. O., Hjorth, J., Ostensen, R., Andersen, M. I., Clasen, J. W., Wucknitz, O., Meylan, G., Magain, P., Stabell, R., Refsdal, S. 1998, *ApJ*, 501, L5
- Blandford, R. & Narayan, R. 1992, *ARA&A*, 30, 311
- Beers, T.C., Flynn, K. & Gebhardt, K., 1990, *AJ*, 100, 32
- Benítez, N., Broadhurst, T., Rosati, P., Courbin, F., Squires, G., Lidman, C., Magain, P., 1999, *ApJ*, 527, 31
- Bernstein, G., Fisher, P., 1999, *AJ*, 118, 14
- Cohen, J.G., Hogg, D.W., Pahre, M.A., Blandford, R.D., Shopbell, P.L., & Richberg, K., 1999, *ApJS*, 120, 171
- Donahue, M., Voit, G. M., Scharf, C. A., Gioia, I. M., Mullis, C. R., Hughes, J. P., Stocke, J. T., 2000, *ApJ*, 527, 525
- Girardi, M., Fadda, D., Giuricin, G., Mardirossian, F., Mezzetti, M., Biviano, A., 1996, *ApJ*, 457, 61
- Hattori, M., Ikebe, Y., Asaoka, I., Takeshima, T., Boehringer, H., Mihara, T., Neumann, D. M., Schindler, S., Tsuru, T., Tamura, T., 1997, *Nature*, 388, 146
- Hjorth, J. et al. 1999, in "Gravitational Lensing: Recent Progress and Future Goals", eds. T. G. Brainerd & C. S. Kochanek, in press.
- Hjorth, J. et al. 2000, *ApJ*, in preparation
- Hjorth, J. & Kneib, J.-P. 2000, *ApJ*, submitted

TABLE 1
JOURNAL OF OBSERVATIONS OF RX J0911+05 DONE AT KECK II USING LRIS ON DECEMBER 7-8,
1999 AND ON MARCH 6-7, 2000.

Filter or λ -range	Exposure Time (sec)	Pixel ($''$) - ($\text{\AA}$)	Seeing FWHM ($''$)
I-band	300	0.215	1.1
R-band	140	0.215	1.1
B-band	200	0.215	1.2
5000–9000 $\text{\AA}$	3000	0.215 - 1.84	1.1
5000–9000 $\text{\AA}$	3000	0.215 - 1.84	0.9

Hoekstra, H., Franx, M., Kuijken, K., 2000, ApJ, 532, 88

Kneib, J.-P., Ellis, R.S., Smail, I., Couch, W., Sharples, R., 1996, 471, 643

Kneib, J.-P., Alloin, D., Mellier, Y., Guilloteau, S., Barvainis, R., Antonucci, R., 1998a, 329, 827

Kneib, J.-P., Alloin, D., Pello, R., 1998b, A&A, 339, L65

Keeton, C. R., & Kochanek, C. S. 1997, ApJ, 487, 42

Kochanek, C. S., Falco, E. E. & Muñoz, J. A. 1999, ApJ, 510, 590

Kochanek, C. S. et al., 2000, astro-ph/9909018

Kundić, T., Cohen, J.G., Blandford R.D. & Lubin, L.M., 1997a, AJ, 114 507

Kundić, T., Hogg, D.W., Blandford, R.D., Cohen, J.G., Lubin, L.M. & Larkin, J.E., 1997b, AJ114, 2276

Kundić, T., Turner, E.L., Colley, W. N., Gott, J. R., Rhoads, J. E., Wang, Y., Bergeron, L., Gloria, K. A., Long, D. C., Malhotra, S., Wambsganss, J., 1997c, ApJ, 482, 75

Kurtz, M.J. & Mink, D.J., 1998, PASP, 110, 934

Landolt, A.U., 1992, AJ, 104, 340

Nolthenius, R., White, S., 1987, MNRAS, 225, 505

Maller, A., Simard, L, Guhathakurta, P., Hjorth, J., Jaunsen, A. O., Flores, R., Primack, J.R., 2000, ApJ, 533, 194

Oke, J. B., Cohen, J. G., Carr, M., Cromer, J., Dingizian, A., Harris, F. H., Labrecque, S., Lucinio, R., Schaal, W., Epps, H., Miller, J., 1995, PASP, 107, 375

Refsdal, S. 1964, MNRAS, 128, 295, 128, 307

Schechter, P. L. et al., 1997, ApJ, 475, L85

Soucail, G., Kneib, J.-P., Jaunsen, A.O, Hjorth, J., Hattori, M., Yamada, T., 2000, A&A, in preparation

Tran, K. H., Kelson, D. D., van Dokkum, P., Franx, M., Illingworth, G. D., Magee, D. 1999, ApJ, 522, 39

This 2-column preprint was prepared with the AAS L^AT_EX macros v5.0.

Fig. 1.— Keck/LRIS BRI image ($4.75' \times 6.15'$ field) showing the central region of the RX J0911+05 cluster. Circled objects were observed in MOS mode, the redshift identification is noted on the side of the circle. The zoomed image corresponds to the HST-NICMOS observation of RX J0911+05 multiple quasar.

TABLE 2
CATALOGUE OF THE CLUSTER GALAXIES IN THE FIELD OF RX J0911+05.

ID	α (J2000)	δ (J2000)	I_{AB}	z	Δz	Line Identification	Cross-correlation template - R	Object type-quality
G1	09:11:26.46	+05:50:14.5	19.99	0.7682	0.0002	H/K/br/G	k0V/g6V - 3.03	A-1
G2	09:11:27.29	+05:50:25.4	20.82	0.7738	0.0003	[O II]/Balmer/H/K/G	spec-em - 3.92	C-1
G3	09:11:27.92	+05:50:49.5	20.90	0.7618	0.0003	H/K/br/G	g6V - 3.43	A-1
G4	09:11:21.17	+05:49:39.8	20.94	0.7693	0.0004	H/K/br/G	gell - 5.40	A-1
G5	09:11:21.40	+05:48:03.8	21.24	0.7697	0.0003	H/K/br/G	gell - 7.50	A-1
G6	09:11:28.36	+05:51:07.9	21.30	0.7650	0.001	wk[O II]/K/H/G	—	C-1
G7	09:11:26.43	+05:50:07.7	21.65	0.7754	0.0005	H/K/br/G	gell - 4.49	A-1
G8	09:11:26.45	+05:50:52.0	21.71	0.776	0.001	wk[O II]/H/K/br	—	C-1
G9	09:11:26.39	+05:49:52.7	21.72	0.7674	0.0003	H/K/br/G	f6V - 3.73	A-1
G10	09:11:22.93	+05:52:40.3	21.87	0.7628	0.0004	MgII/[O II]/H/K	—	C-1
G11	09:11:22.76	+05:48:29.9	21.89	0.7692	0.0002	H/K/br/G	gell - 5.90	A-1
G12	09:11:30.61	+05:50:17.1	21.94	0.766	0.001	H/K/br/G	—	A-2
G13	09:11:31.59	+05:50:37.6	21.98	0.769	0.001	H/K/G	—	A-3
G14	09:11:27.79	+05:50:45.5	22.04	0.760	0.001	H/K/br	—	A-3
G15	09:11:31.22	+05:51:26.6	22.05	0.7745	0.0005	wk[O II]/H δ	—	E-3
G16	09:11:27.82	+05:50:41.2	22.05	0.7758	0.001	wk[O II]/K/H/G	—	C-3
G17	09:11:27.55	+05:51:03.4	22.10	0.7767	0.0003	H/K/br/G	g6III - 4.01	A-1
G18	09:11:24.17	+05:49:35.2	22.16	0.760	0.001	[O II]/H/K/br	—	E-3
G19	09:11:26.78	+05:50:19.0	22.25	0.7640	0.0002	H/K/br/G	g6III - 6.17	A-1
G20	09:11:26.72	+05:50:17.6	22.25	0.7694	0.0006	H/K/br/G	gell - 3.65	A-1
G21	09:11:26.74	+05:50:34.1	22.30	0.7705	0.0005	H/K/br/G	—	A-3
G22	09:11:27.98	+05:49:18.9	22.35	0.770	0.002	H/K/br/G	—	A-3
G23	09:11:30.57	+05:51:13.4	22.42	0.770	0.002	H/K/br/G	—	A-3
L1	09:11:27.56	+05:50:54.21	22.4	0.769	...	...	...	...

NOTE.— R is the cross-correlation parameter. Type and quality classification is similar to the one used by Cohen et al. (1999). L1 is the main galaxy lens.




Fig. 2.— **(left)** Redshift histogram of the redshift survey. **(right)** Redshift histogram of the galaxy distribution at the cluster redshift. The dashed lined corresponds to the mean redshift of the cluster distribution, the dotted line to the standard dispersion of 836^{+180}_{-200} km s⁻¹.




Fig. 3.— **a)** Mass distribution of the cluster (thin lines) overlaid on the K-band image (Courbin, private communication). The thick lines are the critical lines at the redshift $z = 2.80$ of the quasar. **b)** HST/NICMOS-2 F160W archival image of the quadruple quasar RX J0911+05 (PI: Falco). The main lens is composed of two galaxies *L1* and *L2*. The critical lines (thick lines) correspond to the redshift $z = 2.80$ of the quasar. Image A1 is the reference coordinates (0, 0). North is up, East to the left.

This figure "fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0006106v1>